

Sileo®




Dosing

Indication: for the alleviation of acute anxiety and fear associated with noise.

Sileo is administered oromucosally between the dog's cheek and gum at a dose of $125\mu\text{g}/\text{m}^2$

See the table for the dose volume in dots.



| The weight of the dog | Dose/Number of dots |
|-----------------------|--|
| 2.0-5.5 kg | 1 . |
| 5.6-12.0 kg | 2 .. |
| 12.1-20.0 kg | 3 ... |
| 20.1-29.0 kg | 4 |
| 29.1-39.0 kg | 5 <i>(Note: The original image shows 6 dots for this weight range)</i> |
| 39.1-50.0 kg | 6 <i>(Note: The original image shows 7 dots for this weight range)</i> |
| 50.1-62.5 kg | 7 <i>(Note: The original image shows 8 dots for this weight range)</i> |
| 62.6-75.5 kg | 8 <i>(Note: The original image shows 9 dots for this weight range)</i> |
| 75.6-89.0 kg | 9 <i>(Note: The original image shows 10 dots for this weight range)</i> |
| 89.1-100 kg | 10 |

NOTE: If the dose is more than 6 dots, divide the dose between both sides of the mouth.

Administering Sileo®

- **How:** By inserting the syringe tip between the dog's cheek and gum.
- **IMPORTANT: If the gel is swallowed, it may not be effective.**
- **When:** As soon as the dog shows signs of anxiety/fear or when typical triggers of fear are detected or anticipated.
- **No treats or food** within 15 minutes after administration (to allow oromucosal absorption).
- **Can be dosed up to 5 times** during each event, leaving at least 2 hours between each dose.
- **See the Summary of product characteristics before prescribing the medicine.**

Dose into cheek



Not to be swallowed



SILEO® 0.1 mg/ml oromucosal gel for dogs. **ACTIVE SUBSTANCE:** Dexmedetomidine hydrochloride. **INDICATIONS FOR USE:** Alleviation of acute anxiety and fear associated with noise in dogs. **ADMINISTRATION:** Sileo is administered onto the oral mucosa between the dog's cheek and gum at a dose of 125 micrograms/m². **CONTRAINDICATIONS:** Severe cardiovascular disorders, severe systemic disease, hypersensitivity to the active substance or to any of the excipients. Do not use in dogs obviously sedated from previous dosing. **SPECIAL PRECAUTIONS FOR USE:** If the oromucosal gel is swallowed it will become ineffective. Avoid skin, eye or mucosal contact. Wear impermeable disposable gloves. **ADVERSE EVENTS:** Common: transient paleness of mucous membranes at the application site, sedation, emesis and urinary incontinence. See detailed information in the SPC. **LEGAL STATUS:** POM-V. **COMPLETE PRODUCT INFORMATION:** www.ema.europa.eu/en/medicines/veterinary/EPAR/sileo

Dômes Pharma is the trading name of TVM UK Animal Health Ltd. Sileo® is developed and manufactured by Orion Corporation and distributed by TVM-UK Animal Health Ltd, Kirtlington Business Centre, Slade Farm, Kirtlington, Oxfordshire, OX5 3JA, UK. Tel: 0800 0385868 | Email: help@domespharma.com | www.domespharma.co.uk. **Use medicines responsibly.** www.noah.co.uk/responsible. Advice should be sought from the medicine prescriber. Prescription decisions are for the person issuing the prescription alone.